

CLASSIFICATION **CONFIDENTIAL**
 CENTRAL INTELLIGENCE AGENCY
 INFORMATION FROM
 FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT

CD NO.

50X1-HUM

COUNTRY USSR
 SUBJECT Scientific - Medical, antibiotics

DATE OF
 INFORMATION 1950

HOW
 PUBLISHED Monthly periodical

DATE DIST. / *Mar 1951*

WHERE
 PUBLISHED Leningrad

NO. OF PAGES 2

DATE
 PUBLISHED Oct 1950

SUPPLEMENT TO
 REPORT NO.

50X1-HUM

LANGUAGE Russian

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE
 OF THE UNITED STATES WITHIN THE MEANING OF ESPIONAGE ACT 30
 U. S. C. 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION
 OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PRO-
 HIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED.


THIS IS UNEVALUATED INFORMATION

SOURCE Priroda, Vol XXXIX, No 10, p 54.

EFFECT OF PHYTONCIDES ON THE FORMATION OF AGGLUTININS

O. Savchuk

[Since the publication of Professor B. B. Tokin's "Phyton-
 cides -- Essays on Antiseptics of Plant Origin," Press of the
 Academy of Medical Sciences USSR, Moscow, 1948, in-
 creased attention has been paid in the USSR to the study of
 these antibiotics derived from higher plants. Extensive work
 on the subject had been done before. In the course of this work
 it was established that the potent phytoncide contained in gar-
 lic has a glucosidic character (Vestnik Akademii Meditsinskikh
 Nauk SSSR, No 1, 1947). According to Tokin's classification,
 which is schematically outlined below, the tissue juices of on-
 ion, garlic, and of many other plants contain nonexcretory phy-
 tonicides that exert an antibacterial action outside of the plant.
 The active substances in question may be glucosides or, in some
 cases, alkaloids.



- 1 -

CONFIDENTIAL

CLASSIFICATION		CONFIDENTIAL		DISTRIBUTION									
STATE	<input checked="" type="checkbox"/>	NAVY	<input checked="" type="checkbox"/>	NSRB									
ARMY	<input checked="" type="checkbox"/>	AIR	<input checked="" type="checkbox"/>	FBI									

CONFIDENTIAL

CONFIDENTIAL

50X1-HUM

Among the excretory phytoncides, of which some are volatile, the ones having specific action are similar to antibodies, in Tokin's opinion (Mikrobiologiya, Vol XVIII, No 1, 84-90).]

External conditions affect various functions of the living organism. This includes formation of antibodies, which is definitely influenced either positively or negatively by metabolism and various factors having a direct or indirect effect on metabolism. The formation of antibodies can be easily influenced by acting on the digestion or on the nervous system. It is known that a number of substances, when introduced into the body of the immunized animal, stimulate the formation of antibodies.

In our experiments we have studied the effects of garlic, horse-radish, and onion phytoncides on the formation of agglutinins in the body of rabbits which had been immunized with paratyphoid B vaccine. In the course of these experiments, the juice of the plant in question was introduced subcutaneously during 5 days into the body of immunized animals, while another group of immunized animals served as controls.

One to two cubic centimeters of a 5% aqueous solution of the juice of garlic, horse-radish, or onion were injected into the experimental animals twice daily during 5 days. On the 8th day after each immunization, blood was taken from both the experimental and control animals for an agglutination test. As can be seen from the following table, the phytoncides of the three plants used stimulate the formation of agglutinins.

Increases in the Titer of Agglutinins Under the Effect of Phytoncides
of Garlic, Horse-radish, and Onion

<u>Immunization</u>	<u>Titer of Agglutinating Serum of Control Rabbits</u>	<u>Titer of Agglutinating Serum of Rabbits Treated With Phytoncides of:</u>		
		<u>Garlic</u>	<u>Horse-radish</u>	<u>Onion</u>
After first immunization	1:1280-1:2560	1:5120	1:5120	1:2560
After second immunization	1:2560-1:5120	1:10240-1:20480	1:5120-1:10240	1:5120-1:10240
After third immunization	1:5120	1:20480-1:40960	1:20480-1:40960	1:10240

It follows from the results listed in the table that the phytoncides of all three plants are active in stimulating the formation of agglutinins. The juice of garlic is the most effective, while that of onion is the least effective of the three.

- E N D -

- 2 -

CONFIDENTIAL

CONFIDENTIAL